

DRUG STATISTICS SERIES

Number 21

2007 National Drug Strategy Household Survey

State and territory supplement

August 2008

Australian Institute of Health and Welfare

Canberra

Cat. no. PHE 102

© Australian Institute of Health and Welfare 2008

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Media and Communications Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

This publication is part of the Australian Institute of Health and Welfare's Drug statistics series. A complete list of the Institute's publications is available from the Institute's website <www.aihw.gov.au>.

ISBN 978 1 74024 824 2

Suggested citation

Australian Institute of Health and Welfare 2008. 2007 National Drug Strategy Household Survey: State and territory supplement. Drug statistics series no. 21. Cat. no. PHE 102. Canberra: AIHW.

Australian Institute of Health and Welfare

Board Chair
Hon. Peter Collins, AM, QC

Director
Penny Allbon

Any enquiries about or comments on this publication should be directed to:

David Batts
Australian Institute of Health and Welfare
GPO Box 570
Canberra ACT 2601
Phone: (02) 6244 1155
Email: david.batts@aihw.gov.au

Published by the Australian Institute of Health and Welfare

Introduction

About this report

This report presents summary statistics for states and territories from the 2007 National Drug Strategy Household Survey. It supplements the national findings published in the *2007 National Drug Strategy Household Survey: First results* report published in April 2008. No commentary on the results is offered in this supplement.

Some of the results contained in this report have a relative standard error of 50% or greater and have been flagged with an asterisk. They most often arise at low prevalences and from small respondent population. Readers should exercise caution when interpreting such results, especially when making statistical comparisons or statements of trend. Nevertheless, these estimates are valid estimates of the relevant population characteristic and may be quoted as such.

Further, tests of the significance of differences between jurisdictions have not been performed. Readers should not conclude significant differences, even in cases where there are apparently large substantive differences.

Although the survey included respondents aged 12 years or older, only results for those aged 14 years or older are presented in this report, consistent with previous supplements.

About the survey

The survey of around 23,000 Australians living in residential households was the ninth in a series commencing in 1985. The survey covered drug-related attitudes, beliefs and behaviours, as well as support for various drug-related policy options. These topics related to the use of tobacco, alcohol and illicit drugs (including non-medical use of pharmaceuticals).

A full description of the survey, explanatory notes, definitions and a copy of the survey questionnaire can be found in the *First results* publication. A summary of the sample characteristics is at Appendix 1. A *Detailed findings* report is scheduled for release in late 2008. A public-use data set is also available from the Australian Social Science Data Archives at the Australian National University.

Symbols in tables

- Zero
- * Relative standard error greater than 50%
- < 0.1 Non-zero result less than 0.1%

List of tables

Table S1:	Tobacco smoking status: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	3
Table S2:	Daily smoking: proportion of the population aged 14 years or older, by age and sex, states and territories, 2007 (per cent)	4
Table S3:	Alcohol drinking status: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	5
Table S4:	Risk of alcohol-related harm in the long term: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	6
Table S5:	Risk of alcohol-related harm in the short term: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	7
Table S6:	Recent illicit drug use summary: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	8
Table S7:	Recent use of cannabis and any illicit drug excluding cannabis: proportion of the population aged 14 years or older, by age and sex, states and territories, 2007 (per cent)	9
Table S8:	Support for tobacco measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	10
Table S9:	Support for alcohol measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	11
Table S10:	Support for cannabis measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	12
Table S11:	Support for heroin measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)	12
Table SA1:	Sample characteristics, states and territories, 2007	13
Table SA2:	Population estimates, states and territories, as at 30 June 2007	14

Use of tobacco

Table S1: Tobacco smoking status: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Smoking status	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
Daily	18.3	17.9	18.9	15.1	16.8	23.8	14.8	27.9	18.0
Weekly	1.5	1.6	1.4	0.9	1.6	*0.5	1.5	1.5	1.4
Less than weekly	1.5	2.2	1.3	1.4	1.8	2.7	2.0	1.7	1.7
Ex-smoker ^(a)	27.1	26.6	28.7	32.1	28.5	29.5	23.9	25.1	27.9
Never smoked ^(b)	51.7	51.6	49.7	50.5	51.3	43.4	57.8	43.9	50.9
Females									
Daily	14.5	15.2	15.4	14.5	16.2	21.6	14.6	22.5	15.2
Weekly	1.0	1.3	1.4	1.5	1.3	*0.7	*0.3	*0.8	1.2
Less than weekly	1.2	1.2	1.2	1.5	1.7	*0.5	1.3	*0.8	1.3
Ex-smoker ^(a)	22.3	22.2	22.7	24.5	19.9	23.7	25.7	19.6	22.4
Never smoked ^(b)	60.9	60.1	59.3	58.1	60.9	53.6	58.1	56.3	59.8
Persons									
Daily	16.4	16.5	17.2	14.8	16.5	22.7	14.7	25.3	16.6
Weekly	1.2	1.5	1.4	1.2	1.5	0.6	0.9	1.2	1.3
Less than weekly	1.4	1.7	1.3	1.4	1.8	1.6	1.6	1.3	1.5
Ex-smoker ^(a)	24.7	24.4	25.7	28.3	24.1	26.5	24.8	22.4	25.1
Never smoked ^(b)	56.3	55.9	54.5	54.3	56.2	48.6	57.9	49.8	55.4

(a) Smoked at least 100 cigarettes (or equivalent amount of tobacco) in lifetime, but no longer smokes.

(b) Never smoked more than 100 cigarettes (or equivalent amount of tobacco) in lifetime.

* Relative standard error greater than 50%

Table S2: Daily smoking: proportion of the population aged 14 years or older, by age and sex, states and territories, 2007 (per cent)

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
14–19	6.6	5.4	5.7	*5.2	*5.5	*3.0	*9.2	21.0	6.0
20–29	22.6	24.5	28.3	19.2	16.5	31.1	21.4	36.1	23.7
30–39	24.6	21.5	23.0	19.6	18.0	37.0	20.9	24.8	22.7
40–49	21.3	22.7	22.5	19.2	21.4	32.8	10.8	26.4	21.8
50–59	21.0	19.6	20.0	13.2	25.6	24.7	11.1	32.4	20.1
60+	10.9	10.4	10.3	10.3	11.5	13.0	11.2	22.9	10.8
Aged 14+	18.3	17.9	18.9	15.1	16.8	23.8	14.8	27.9	18.0
Females									
14–19	7.1	8.5	13.1	6.1	*5.1	13.5	*11.2	*5.6	8.7
20–29	16.5	21.1	17.5	20.6	20.1	29.6	25.1	22.0	19.0
30–39	18.9	18.0	16.9	18.9	23.5	32.4	13.6	29.1	18.9
40–49	19.3	20.8	21.6	14.4	26.7	34.8	16.1	27.4	20.6
50–59	15.7	13.7	14.1	18.1	12.5	16.6	16.7	22.5	15.0
60+	8.9	8.4	9.8	7.6	9.2	8.8	*3.7	19.1	8.8
Aged 14+	14.5	15.2	15.4	14.5	16.2	21.6	14.6	22.5	15.2
Persons									
14–19	6.9	6.9	9.3	5.7	5.3	8.4	10.2	13.6	7.3
20–29	19.6	22.8	23.0	19.9	18.2	30.4	23.2	29.2	21.4
30–39	21.8	19.7	19.9	19.3	20.7	34.6	17.2	26.9	20.8
40–49	20.3	21.7	22.1	16.8	24.1	33.8	13.5	26.9	21.2
50–59	18.3	16.6	17.1	15.6	18.9	20.6	14.0	27.8	17.5
60+	9.8	9.3	10.0	8.9	10.3	10.8	7.2	21.2	9.7
Aged 14+	16.4	16.5	17.2	14.8	16.5	22.7	14.7	25.3	16.6

* Relative standard error greater than 50%

Use of alcohol

Table S3: Alcohol drinking status: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Drinking status	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
Daily	10.7	10.2	10.9	12.8	11.0	9.0	7.8	14.8	10.8
Weekly	45.0	45.9	47.4	51.8	47.6	49.6	52.8	49.8	46.8
Less than weekly	28.0	29.6	28.6	24.9	28.9	30.3	28.1	23.7	28.3
Ex-drinker ^(a)	5.9	5.6	6.5	4.5	6.4	4.5	6.3	6.7	5.8
Never a full serve of alcohol	10.3	8.7	6.5	5.9	6.2	6.7	5.1	5.0	8.2
Females									
Daily	5.9	4.6	5.7	6.9	5.0	4.8	5.6	6.2	5.5
Weekly	34.2	34.4	37.1	42.1	36.2	31.9	43.5	37.9	35.9
Less than weekly	36.0	41.0	39.9	34.1	40.1	45.9	38.7	38.4	38.5
Ex-drinker ^(a)	8.9	8.0	7.4	7.2	8.5	7.7	3.2	9.5	8.1
Never a full serve of alcohol	14.9	12.1	9.8	9.6	10.3	9.6	9.0	7.9	12.1
Persons									
Daily	8.3	7.3	8.3	9.8	7.9	6.8	6.6	10.7	8.1
Weekly	39.5	40.0	42.2	46.9	41.8	40.5	48.0	44.1	41.3
Less than weekly	32.1	35.4	34.3	29.5	34.6	38.3	33.5	30.7	33.5
Ex-drinker ^(a)	7.5	6.8	7.0	5.9	7.5	6.2	4.7	8.0	7.0
Never a full serve of alcohol	12.7	10.4	8.2	7.8	8.2	8.2	7.1	6.4	10.1

(a) Consumed at least a full serve of alcohol, but not in the past 12 months.

Table S4: Risk of alcohol-related harm in the long term: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Risk status	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
Abstainers ^(a)	16.2	14.3	13.0	10.5	12.5	11.1	11.3	11.7	14.0
Low risk	73.8	76.7	75.5	78.8	77.9	76.1	80.1	71.0	75.8
Risky or high risk ^(b)									
Risky	6.4	5.4	6.7	7.0	5.5	7.2	5.2	9.3	6.2
High risk	3.5	3.7	4.8	3.7	4.0	5.6	3.3	8.0	3.9
<i>Total risky or high risk</i>	<i>10.0</i>	<i>9.1</i>	<i>11.5</i>	<i>10.7</i>	<i>9.5</i>	<i>12.7</i>	<i>8.6</i>	<i>17.3</i>	<i>10.2</i>
Females									
Abstainers ^(a)	23.9	20.1	17.2	16.9	18.8	17.4	12.2	17.4	20.1
Low risk	66.1	70.8	70.7	70.8	72.3	71.4	76.6	67.0	69.4
Risky or high risk ^(b)									
Risky	7.5	6.6	8.9	9.0	6.6	6.5	7.9	10.8	7.6
High risk	2.6	2.5	3.2	3.3	2.3	4.7	3.2	4.8	2.8
<i>Total risky or high risk</i>	<i>10.1</i>	<i>9.1</i>	<i>12.1</i>	<i>12.3</i>	<i>8.9</i>	<i>11.2</i>	<i>11.1</i>	<i>15.6</i>	<i>10.5</i>
Persons									
Abstainers ^(a)	20.1	17.2	15.1	13.7	15.7	14.4	11.8	14.4	17.1
Low risk	69.9	73.7	73.1	74.8	75.1	73.7	78.3	69.1	72.6
Risky or high risk ^(b)									
Risky	6.9	6.0	7.8	8.0	6.0	6.8	6.6	10.0	6.9
High risk	3.1	3.1	4.0	3.5	3.2	5.1	3.3	6.5	3.4
<i>Total risky or high risk</i>	<i>10.0</i>	<i>9.1</i>	<i>11.8</i>	<i>11.5</i>	<i>9.2</i>	<i>12.0</i>	<i>9.9</i>	<i>16.5</i>	<i>10.3</i>

(a) Not consumed alcohol in the past 12 months.

(b) For males, consumption of 29 or more standard drinks per week; for females, consumption of 15 or more standard drinks per week.
A standard drink is 10 grams (or 12.5 millilitres) of pure alcohol.

Table S5: Risk of alcohol-related harm in the short term: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Risk status	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
Abstainers ^(a)	16.2	14.3	13.0	10.5	12.5	11.1	11.3	11.7	14.0
Low risk	48.5	46.6	46.0	48.2	48.2	40.5	50.6	35.9	47.2
Risky or high risk ^(b)									
At least yearly	13.5	16.2	15.2	16.1	14.6	20.9	13.9	19.6	15.1
At least monthly	12.8	13.4	15.7	16.3	16.3	15.7	15.0	18.6	14.3
At least weekly	8.9	9.5	10.0	8.9	8.4	11.8	9.1	14.2	9.3
<i>Total risky or high risk</i>	<i>35.2</i>	<i>39.1</i>	<i>40.9</i>	<i>41.3</i>	<i>39.2</i>	<i>48.4</i>	<i>38.0</i>	<i>52.4</i>	<i>38.7</i>
Females									
Abstainers ^(a)	23.9	20.1	17.2	16.9	18.8	17.4	12.2	17.4	20.1
Low risk	47.6	50.9	49.3	50.1	49.5	51.4	53.5	45.0	49.3
Risky or high risk ^(b)									
At least yearly	12.8	13.0	15.2	12.6	13.1	12.3	16.1	14.0	13.4
At least monthly	9.8	10.2	11.8	13.1	13.0	11.1	11.7	13.8	10.9
At least weekly	5.9	5.8	6.5	7.3	5.7	7.8	6.5	9.8	6.2
<i>Total risky or high risk</i>	<i>28.5</i>	<i>29.0</i>	<i>33.5</i>	<i>33.0</i>	<i>31.7</i>	<i>31.2</i>	<i>34.3</i>	<i>37.6</i>	<i>30.5</i>
Persons									
Abstainers ^(a)	20.1	17.2	15.1	13.7	15.7	14.4	11.8	14.4	17.1
Low risk	48.1	48.8	47.7	49.2	48.9	46.1	52.1	40.2	48.3
Risky or high risk ^(b)									
At least yearly	13.1	14.5	15.2	14.4	13.8	16.5	15.0	16.9	14.2
At least monthly	11.3	11.8	13.7	14.7	14.6	13.3	13.3	16.3	12.6
At least weekly	7.4	7.6	8.3	8.1	7.0	9.7	7.8	12.1	7.8
<i>Total risky or high risk</i>	<i>31.8</i>	<i>33.9</i>	<i>37.2</i>	<i>37.1</i>	<i>35.4</i>	<i>39.6</i>	<i>36.1</i>	<i>45.3</i>	<i>34.6</i>

(a) Not consumed alcohol in the past 12 months.

(b) For males, consumption of 7 or more standard drinks on one occasion; for females, consumption of 5 or more standard drinks on one occasion. A standard drink is 10 grams (or 12.5 millilitres) of pure alcohol.

Use of illicit drugs

Table S6: Recent^(a) illicit drug use summary: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Substance/behaviour	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Marijuana/cannabis	8.0	8.8	9.5	10.8	10.2	10.8	9.1	13.8	9.1
Pain-killers/analgesics ^(b)	2.2	2.7	2.8	2.5	2.8	3.0	2.0	3.3	2.5
Tranquillisers/sleeping pills ^(b)	1.2	1.5	1.4	1.6	1.2	1.0	0.9	2.1	1.4
Steroids ^(b)	0.1	*0.1	0.2	*0.1	*<0.1	—	*0.1	—	0.1
Barbiturates ^(b)	0.1	*0.1	0.2	*0.1	—	*0.3	*0.2	*0.1	0.1
Inhalants	0.4	0.5	0.5	0.3	*0.1	0.6	0.6	*0.1	0.4
Heroin	0.2	0.3	0.2	*0.2	*0.1	*0.3	—	*0.3	0.2
Methadone or Buprenorphine ^(c)	*< 0.1	*< 0.1	*0.1	*0.1	—	*0.2	*0.1	*0.2	0.1
Other opiates/opioids ^(b)	0.2	*0.1	0.3	0.3	*0.1	*0.4	*0.1	*0.3	0.2
Meth/amphetamine ^(b)	1.8	2.3	2.0	4.2	2.6	1.7	2.3	2.3	2.3
Cocaine	2.0	1.6	1.4	1.8	1.3	0.8	1.4	0.9	1.6
Hallucinogens	0.5	0.5	0.6	1.0	0.9	1.0	0.8	0.9	0.6
Ecstasy	3.4	3.6	3.7	4.1	2.9	2.4	4.7	4.2	3.5
Ketamine	0.3	*0.1	*0.1	*0.1	*0.3	*0.3	*0.2	—	0.2
GHB	0.2	*0.1	*< 0.1	*0.1	*0.1	—	*< 0.1	*0.1	0.1
Injected drugs	0.4	0.5	0.4	0.7	0.3	0.5	0.5	0.5	0.5
<i>Any illicit</i>	<i>12.1</i>	<i>12.8</i>	<i>13.7</i>	<i>16.2</i>	<i>14.7</i>	<i>14.8</i>	<i>13.8</i>	<i>20.4</i>	<i>13.4</i>

(a) Used in the past 12 months.

(b) For non-medical purposes.

(c) Non-maintenance.

* Relative standard error greater than 50%

Table S7: Recent^(a) use of cannabis and any illicit drug excluding cannabis^(b): proportion of the population aged 14 years or older, by age and sex, states and territories, 2007 (per cent)

Drug group and age	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Males									
Cannabis									
14–24	15.8	20.2	20.2	25.0	16.0	22.4	18.6	28.9	19.0
25–39	17.8	19.3	17.1	20.9	23.5	26.6	18.7	23.5	19.0
40+	5.8	4.4	5.5	5.3	7.9	7.3	3.5	9.9	5.5
<i>All ages</i>	10.7	11.3	11.5	13.3	13.1	14.5	11.0	18.1	11.6
Any illicit excluding cannabis									
14–24	3.3	*1.6	4.6	*2.2	*1.3	*2.5	9.0	11.9	3.1
25–39	5.5	4.6	5.5	5.9	5.9	*2.0	*3.9	4.7	5.2
40+	3.3	3.3	2.8	4.2	3.6	3.1	*1.6	5.4	3.3
<i>All ages</i>	3.8	3.3	3.9	4.3	3.7	2.7	3.8	6.4	3.8
Females									
Cannabis									
14–24	10.2	15.4	18.5	19.3	19.2	14.4	16.7	9.9	15.0
25–39	8.6	10.7	10.7	11.4	9.8	15.5	11.1	10.7	10.1
40+	2.4	1.6	2.3	3.3	3.0	2.1	2.1	7.8	2.3
<i>All ages</i>	5.3	6.4	7.5	8.4	7.4	7.3	7.4	9.2	6.6
Any illicit excluding cannabis									
14–24	4.9	4.3	4.5	8.8	8.2	*5.9	*6.1	8.7	5.4
25–39	4.7	4.9	4.8	7.8	6.0	*2.4	4.1	7.5	5.1
40+	3.1	3.3	3.5	3.7	3.1	5.0	4.7	3.0	3.4
<i>All ages</i>	3.8	3.9	4.0	5.7	4.6	4.6	4.8	5.8	4.2
Persons									
Cannabis									
14–24	13.1	17.9	19.4	22.1	17.5	18.3	17.6	19.7	17.0
25–39	13.0	14.9	13.9	16.3	16.6	21.0	15.0	17.1	14.5
40+	4.0	2.9	3.9	4.3	5.4	4.6	2.8	8.9	3.9
<i>All ages</i>	8.0	8.8	9.5	10.8	10.2	10.8	9.1	13.8	9.1
Any illicit excluding cannabis									
14–24	4.1	2.9	4.6	5.5	4.6	4.2	7.5	10.3	4.2
25–39	5.1	4.7	5.1	6.8	5.9	2.2	4.0	6.1	5.2
40+	3.2	3.3	3.2	4.0	3.3	4.1	3.3	4.3	3.3
<i>All ages</i>	3.8	3.6	3.9	5.0	4.2	3.7	4.3	6.1	4.0

(a) Used in the past 12 months.

(b) Excludes those that have used cannabis in the past 12 months, whether or not they had also used other drugs.

* Relative standard error greater than 50%

Community attitudes

Tobacco

Table S8: Support^(a) for tobacco measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Measure	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Banning smoking in the workplace	81.8	82.7	80.6	83.6	84.0	79.4	84.4	69.6	82.0
Banning smoking in pubs/clubs	76.4	78.2	75.9	80.0	75.0	78.1	79.7	62.2	77.0
Increasing tax on tobacco products to pay for health education	67.1	67.6	66.4	70.5	64.5	61.8	70.1	60.8	67.1
Increasing tax on tobacco products to contribute to treatment costs	68.9	68.9	67.4	72.5	64.9	64.2	72.3	62.9	68.6
Increasing tax on tobacco products to discourage smoking	66.0	65.4	64.4	70.9	64.1	60.2	67.5	58.2	65.7
Making it harder to buy tobacco in shops	65.7	66.4	65.6	70.8	67.2	64.9	65.6	57.8	66.4
Stricter enforcement of law against supplying to minors	89.7	89.9	90.6	91.2	90.1	90.8	89.1	84.2	90.1
Stricter penalties for sale or supply of tobacco products to minors	87.4	87.7	87.6	88.4	86.8	89.1	86.3	82.6	87.5
Bans on point of sale advertising and display of tobacco products	74.4	73.6	71.0	77.8	73.2	73.5	71.5	64.9	73.6
Implementing a licensing scheme for tobacco retailers	71.4	72.2	70.6	73.8	70.5	72.9	71.0	63.9	71.6

(a) Support or strongly support; all results as a proportion of those that expressed an opinion.

Alcohol

Table S9: Support^(a) for alcohol measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Measure	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Increasing the price of alcohol	26.3	24.3	22.0	23.2	21.8	22.5	24.0	21.7	24.1
Reducing the number of outlets that sell alcohol	33.6	34.4	30.8	26.9	28.7	31.4	28.9	43.2	32.2
Reducing trading hours for pubs and clubs	44.0	41.4	35.9	28.1	33.3	34.7	34.8	37.0	38.9
Raising the legal drinking age	47.1	44.1	48.6	47.0	47.3	44.7	34.1	38.9	46.3
Increasing the number of alcohol-free public events	63.9	63.5	61.5	62.4	58.0	58.7	62.4	54.2	62.5
Increasing the number of alcohol-free dry zones	66.0	61.4	60.2	62.4	64.6	59.9	62.1	56.4	63.0
Serving only low-alcohol beverages at sporting events	62.4	59.8	60.6	59.2	52.8	57.4	57.3	55.5	60.1
Limiting TV advertising until after 9:30 p.m.	73.2	72.5	69.9	73.8	71.4	74.2	69.6	64.1	72.2
Banning alcohol sponsorship of sporting events	51.9	48.0	42.8	53.6	46.3	45.5	49.3	36.3	48.5
More severe penalties for drink driving	86.6	86.6	86.8	85.0	85.0	87.3	84.8	84.2	86.3
Stricter laws against serving drunk customers	84.2	83.4	82.8	82.5	83.1	82.7	82.2	80.1	83.3
Restricting late night trading of alcohol	61.9	59.1	57.0	50.6	52.3	54.8	52.5	53.9	58.0
Strict monitoring of late night licensed premises	76.6	77.0	74.7	71.2	70.5	76.2	71.9	69.8	75.2
Increasing the size of standard drink labels on alcohol containers	66.2	67.0	66.5	63.7	62.7	63.1	63.7	58.5	65.8
Requiring national drinking guidelines on alcohol containers	71.6	70.6	71.1	71.3	70.6	67.5	69.7	59.9	70.9
Increasing tax on alcohol to pay for health, education and treatment of alcohol-related problems	43.5	41.4	38.5	41.8	38.4	39.9	42.6	35.2	41.3

(a) Support or strongly support; all results as a proportion of those that expressed an opinion.

Illicit drugs

Table S10: Support^(a) for cannabis measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Measure	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
A change in legislation permitting the use of marijuana for medical purposes	68.4	68.1	69.0	69.3	71.4	70.0	72.8	71.0	68.6
A clinical trial for people to use marijuana to treat medical conditions	73.0	73.1	73.7	73.1	75.2	74.3	78.1	75.1	73.6

(a) Support or strongly support; all results as a proportion of those that expressed an opinion.

Table S11: Support^(a) for heroin measures: proportion of the population aged 14 years or older, states and territories, 2007 (per cent)

Measure	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
Needle and syringe programs	63.8	67.3	68.4	72.8	67.2	67.8	71.1	64.5	67.0
Methadone maintenance programs	63.9	69.1	68.7	73.2	68.6	72.9	69.8	61.8	67.7
Treatment with drugs other than methadone	67.2	68.5	69.2	72.6	66.1	72.2	69.2	64.0	68.5
Regulated injecting rooms	49.8	50.4	47.8	52.4	49.0	54.1	56.2	45.9	49.9
Trial of prescribed heroin	31.9	33.5	31.1	37.1	32.7	34.8	38.4	30.0	32.9
Rapid detoxification therapy	78.2	77.9	79.7	83.4	75.7	82.8	75.6	74.6	78.8
Use of Naltrexone	74.0	72.9	75.8	81.9	69.9	78.5	76.1	69.7	74.7

(a) Support or strongly support; all results as a proportion of those that expressed an opinion.

Appendix 1: Sample characteristics

Table SA1: Sample characteristics, states and territories, 2007

Characteristic	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Aus
	(number)								
Sample size (14+)									
Drop-and-collect	5,505	4,060	3,569	1,977	1,594	944	978	816	19,443
CATI	1,006	674	586	394	353	199	61	196	3,469
<i>Total</i>	<i>6,511</i>	<i>4,734</i>	<i>4,155</i>	<i>2,371</i>	<i>1,947</i>	<i>1,143</i>	<i>1,039</i>	<i>1,012</i>	<i>22,912</i>
	(per cent)								
Completion rate^(a)									
Drop-and-collect	51	48	59	58	58	61	65	55	54
CATI	39	38	37	42	39	41	42	44	39
<i>Total</i>	<i>49</i>	<i>47</i>	<i>56</i>	<i>55</i>	<i>55</i>	<i>58</i>	<i>64</i>	<i>53</i>	<i>52</i>
	(number)								
Average design effect^(b)									
Drop-and-collect	1.59	1.60	1.73	1.31	1.30	1.40	1.42	1.36	1.71
CATI	1.03	0.99	1.00	1.00	0.99	0.99	1.00	0.99	1.12
<i>Total</i>	<i>1.58</i>	<i>1.65</i>	<i>1.71</i>	<i>1.34</i>	<i>1.31</i>	<i>1.42</i>	<i>1.37</i>	<i>1.33</i>	<i>1.72</i>

(a) This represents a mixture of response rate and participation rate across the two survey modes.

(b) The design effect is the ratio of the variance estimate from this sample compared with that from a simple random sample of the same size. It indicates the degree of under-estimation of variance that may arise from not taking into account the complex nature of the survey.

Appendix 2: Population estimates

Table SA2: Population estimates, states and territories, as at 30 June 2007

Age group	NSW	Vic	Qld	WA	SA	Tas	ACT	NT	Total
Males									
14–19	285,800	215,900	180,400	92,100	65,200	21,000	14,800	10,200	885,400
20–29	479,800	375,400	296,100	153,500	106,800	29,600	29,700	18,300	1,489,400
30–39	492,100	380,900	299,400	155,100	107,100	30,700	26,100	18,400	1,510,100
40–49	491,100	371,500	298,200	157,400	114,700	35,000	24,100	16,600	1,508,900
50–59	429,000	319,500	264,100	137,500	103,500	33,700	21,000	13,300	1,321,600
60+	597,500	443,200	346,400	168,400	148,700	47,200	22,700	10,000	1,784,200
14–24	526,600	406,600	332,200	171,600	121,400	36,900	30,300	19,400	1,645,200
25–39	731,200	565,600	443,700	229,200	157,600	44,400	40,300	27,500	2,239,800
40+	1,517,600	1,134,200	908,600	463,300	366,900	115,900	67,700	39,800	4,614,800
Aged 14+	2,775,400	2,106,400	1,684,500	864,000	646,000	197,200	138,400	86,700	8,499,700
Females									
14–19	270,900	205,300	171,500	86,500	61,500	19,900	14,100	9,300	839,100
20–29	472,400	366,100	288,800	143,400	102,600	29,600	28,700	17,500	1,449,200
30–39	500,900	387,700	302,400	150,200	105,900	32,000	26,200	17,600	1,523,100
40–49	499,400	380,000	305,300	155,300	116,000	36,400	25,200	15,300	1,533,000
50–59	433,300	328,000	263,800	134,400	106,800	34,200	22,100	11,800	1,334,600
60+	694,700	518,200	379,600	185,200	178,700	54,400	26,100	8,300	2,045,300
14–24	506,100	390,000	320,000	160,200	115,600	35,300	28,500	17,700	1,573,500
25–39	738,100	569,100	442,700	219,800	154,500	46,200	40,500	26,800	2,237,900
40+	1,627,400	1,226,300	948,700	474,900	401,400	125,000	73,500	35,400	4,912,900
Aged 14+	2,871,600	2,185,400	1,711,400	855,000	671,400	206,600	142,400	79,800	8,724,300
Persons									
14–19	556,800	421,200	351,900	178,600	126,700	40,900	28,900	19,500	1,724,600
20–29	952,200	741,500	584,900	296,900	209,400	59,200	58,400	35,800	2,938,600
30–39	993,000	768,700	601,800	305,300	213,000	62,700	52,300	36,100	3,033,200
40–49	990,500	751,500	603,500	312,700	230,700	71,400	49,300	31,900	3,041,900
50–59	862,300	647,500	527,800	272,000	210,200	67,900	43,100	25,100	2,656,200
60+	1,292,200	961,400	726,000	353,500	327,400	101,600	48,800	18,300	3,829,500
14–24	1,032,700	796,600	652,200	331,800	237,000	72,200	58,800	37,000	3,218,700
25–39	1,469,300	1,134,700	886,400	449,000	312,100	90,600	80,800	54,300	4,477,700
40+	3,145,000	2,360,400	1,857,300	938,200	768,300	240,900	141,200	75,200	9,527,700
Aged 14+	5,647,000	4,291,800	3,395,900	1,719,000	1,317,400	403,700	280,800	166,500	17,224,000

Source: AIHW population database, as at June 2008. Population estimates updated since publication of *First results*.